



## **Important Correction Notice** ***Handbook on Injectable Drugs, 19<sup>th</sup> edition***

ASHP wishes to inform you of a correction to our publication, the *Handbook on Injectable Drugs, 19<sup>th</sup> edition*. This correction notice applies only to the printed edition.

On page 224 in the cefepime hydrochloride monograph, reconstitution instructions for the intramuscular injection of cefepime appear as "...reconstitute the 1- and 2-g vials with 1.3 and 2.4 mL, respectively..."

*The corrected version of the paragraph is as follows:*

For intramuscular administration, reconstitute the **500-mg** and **1-g** vials with 1.3 and 2.4 mL, respectively, of sterile water for injection, sodium chloride 0.9%, dextrose 5%, lidocaine hydrochloride 0.5 or 1%, or bacteriostatic water for injection preserved with parabens or benzyl alcohol to yield a solution with a cefepime concentration of 280 mg/mL.<sup>3295</sup>

We urge you to correct this error in all copies of the *Handbook on Injectable Drugs, 19<sup>th</sup> edition* immediately and to communicate it to others on your staff who may use the book. The Interactive *Handbook on Injectable Drugs* has been corrected.

***For print copy purchasers of the Handbook on Injectable Drugs:***

**If you have a print copy of the *Handbook on Injectable Drugs, 19<sup>th</sup> edition*, please keep this correction notice with your copy and inform all others who may refer to the book.**

***For purchasers of the Interactive Handbook on Injectable Drugs:***

**Please note that the correction has already been made.**